

(12)

G

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : A61K 31/47, 37/00	A1	(11) International Publication Number: WO 91/19498 (43) International Publication Date: 26 December 1991 (26.12.91)
(21) International Application Number: PCT/US91/03788 (22) International Filing Date: 5 June 1991 (05.06.91) (30) Priority data: 535,672 11 June 1990 (11.06.90) US (71) Applicant: THE DU PONT MERCK PHARMACEUTICAL COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: ACKERMAN, Neil, Richard ; 807 Marina View, Alameda, CA 94501 (US). JAFFEE, Bruce, Donald ; 3206 Heathwood Road, Wilmington, DE 19810 (US). (74) Agents: FERGUSON, Blair, Q. et al.; The Du Pont Merck Pharmaceutical Company, Legal/Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).	(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>	
(54) Title: TREATMENT OF ORGAN TRANSPLANTATION REJECTION (57) Abstract 4-Quinoline-carboxylic acid derivatives, such as 2-(2'-fluoro-1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid, in combination with other immunosuppressive agents are useful for the treatment and/or prevention of transplantation rejection, graft vs. host disease, autoimmune diseases, and chronic inflammatory disease.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LJ	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

Title

Treatment of Organ Transplantation Rejection

Background of the Invention

5 U.S. Patent 4,680,299 (Hesson), granted July 14, 1987, describes 4-quinoline-carboxylic acid derivatives as tumor inhibiting agents. Commonly assigned European Patent Application Number 89107099.7, published 11/2/89 describes 4-quinoline-carboxylic acid derivatives as
10 immunomodulatory and anti-inflammatory agents.

U.S. Patent 4,847,381 (Sutherland et al.) teaches substituted 4-quinoline-carboxylic acid compounds useful for the treatment of arthritis and useful as immunosuppressive agents.

15 Presently, cyclosporin A, an immunosuppressive agent, used in combination with other adjunctive therapies, such as azathioprine (AZA) and corticosteroids, is the treatment of choice for the prevention of organ transplantation rejection. Other
20 immunosuppressive agents, including but not limited to azathioprine, corticosteroids (such as prednisone), OKT3, FK506, mycophenolic acid or the morpholinethylester thereof, 15-dioxyspergualin, rapamycin, mizoribine, misoprostol and anti-
25 interleukin-2 (IL-2) receptor antibodies, have been used or are suggested to be useful in the treatment and/or prevention of organ transplantation rejection.


Use of any of these known immunosuppressive compounds, either alone or in combination, is associated
30 with a high incidence of side effects such as nephrotoxicity and/or hepatotoxicity. The 4-quinoline-carboxylic acid compounds useful in the present invention have a unique mechanism of action compared to other known immunosuppressive agents, and therefore have
35 not been associated with the nephrotoxicity and hepatotoxicity seen with other immunosuppressive agents

such as cyclosporin A and AZA. In addition, the combination of a 4-quinoline-carboxylic acid with known immunosuppressive agents has a synergistic effect in terms of inhibition of inflammation in animal models.

- 5 This synergistic effect is seen at suboptimal doses of each immunosuppressive agent, thus, suggesting that known immunosuppressive agents could be used in combination with a 4-quinoline-carboxylic acid compound, each at lower doses with an associated lower incidence
10 of side effects.

Summary of the Invention



- According to the present invention, there are provided methods of treating and/or preventing organ transplantation rejection, graft versus host disease,
15 psoriasis and autoimmune diseases, including but not limited to rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, myasthenia gravis as well as chronic inflammatory disease including but not limited to Chron's disease and primary billiary
20 cirrhosis, in a mammal, said methods comprising administering to the mammal in an effective amount for the treatment of a desired aforesaid disease a combination of: (a) at least one immunosuppressive agent preferably selected from the group consisting of
25 cyclosporin A, azathioprine, a corticosteroid such as prednisone or prednisolone, OKT3, FK506, mycophenolic acid or the morpholinethylester thereof, 15-dioxyspergualin, rapamycin, mizoribine, misoprostol and anti-interleukin-2 (IL-2) receptor antibodies, and (b) a
30 compound of the formula:



or a pharmaceutically acceptable salt thereof
wherein:


5

R is




R¹ is CH₃CH₂(CH₃)CH, alkyl of 5 to 12 carbon atoms,
cyclohexyl,

10



R¹ can be in addition alkyl of 3 to 4 carbon atoms;

15 R² is

R^3 is H, alkoxy of 1 to 3 carbon atoms, alkyl of 1 to 2 carbon atoms, hydroxy or alkanoyloxy of 2 to 6 carbon atoms;

5 when R^3 is hydroxy or alkanoyloxy of 2 to 6 carbon atoms, R^1 can be in addition, trifluoromethyl, halogen, hydroxy, alkyl of 1 to 6 carbon atoms and cycloalkyl of 1 to 6 carbon atoms;

R^4 is CO_2H or CO_2R^{11} ;

10 R^5 , R^6 , R^7 and R^8 are independently H, F, Cl, Br, I, CH_3 , CF_3 , SCH_3 or alkyl of 1 to 3 carbon atoms, at least two of R^5 , R^6 , R^7 and R^8 being H;

R^9 and R^{9A} are independently H or alkyl of 1 to 3 carbon atoms;

15 R^{11} is $(CH_2)_2-4NR^9R^{9A}$;

W, Y and Z are independently H, F, Cl, Br, alkyl of 1 to 5 carbon atoms, NO_2 , OH, CF_3 or OCH_3 ; and

m is 0 or 1;

with the following provisos:

- 20 (1) R^5 , R^6 and R^7 cannot all be H;
(2) when R^4 is $CO_2CH_2CH_2N(CH_3)_2$, R^6 is CH_2CH_3 , or R^7 is Cl, R^1 cannot be cyclohexyl;
(3) when R^1 is cyclohexyl and R^3 is H,
25 R^6 must be Cl or F, but R^6 and R^8 cannot both be Cl; and
(4) when R^6 is CH_3 , then R^7 cannot be Cl.

30 Current recommended therapy for the prevention of organ transplantation rejection and related disorders, including graft versus host disease, traditionally involves patient treatment with cyclosporin A and adjunctive therapy with corticosteroids and other
35 immunosuppressive drugs (Immune Modulation Agents and

Their Mechanisms, ISBN 0-8247-7178-8, 1984, pp 191-228, Jacobs and Elgin "Cyclosporin A, Current Status, Including the Cape Town Experience" Transplantation and Clinical Immunology, Volume XX Combined Immuno-


- 5 suppressive Therapy in Transplantation ISBN 0-444-81068-4, 1989). Our results indicate that 2-phenyl-4-quinoline-carboxylic acids and their derivatives will be useful as agents to be used in combination with compounds currently used in these
10 clinical regimens.

The significant clinically observed toxicities associated with cyclosporin A (nephrotoxicity) and AZA (hepatotoxicity), have not been seen with 2-phenyl-4-quinoline-carboxylic acids and their derivatives.

- 15 Moreover, 2-phenyl-4-quinoline-carboxylic acids and their derivatives have a unique mechanism of action (inhibition of dihydroorotate dehydrogenase) which is distinct from other available immunosuppressive agents. Our results show that 2-phenyl-4-quinoline-carboxylic
20 acids and their derivatives have synergistic activity when included in the currently accepted regimen of drug therapy for the prevention of organ transplantation rejection and related complications. More particularly, these 2-phenyl-4-quinoline-carboxylic acid derivatives
25 should be useful at reduced doses of both the quinoline-carboxylic acid compound and other immunosuppressive agents used in combination therewith.

Preferred Embodiments

- 30 Preferred compounds useful in the present invention have the formula:




or a pharmaceutically acceptable salt thereof
wherein:

- 5 R^1 is cyclohexyl; phenyl; phenyl substituted with one halogen; alkyl of 1 to 5 carbon atoms or CF_3 ; phenoxy; or phenoxy substituted with one halogen or alkyl of 1 to 5 carbon atoms; and/or
- R^3 is H or alkyl of 1 to 2 carbon atoms; and/or
- 10 R^4 is CO_2H , a sodium or potassium salt thereof; or CO_2R^{11} ; and/or
- R^5 and R^6 are independently H, halogen, CH_3 or CF_3 ; and/or
- R^7 and R^8 are independently H or halogen; and/or
- 15 R^{11} is $(CH_2)_2-4NR^9R^{9A}$; and/or
- R^9 and R^{9A} are independently alkyl of 1 to 3 carbon atoms;
- provided that R^5 , R^6 and R^7 cannot all be H; when R^1 is cyclohexyl and R^3 is H, R^6 must be Cl or F, but R^6 and R^8 cannot both be Cl; when R^6 is CH_3 , then R^7 cannot be Cl; and when R^4 is CO_2H , R^1 or R^2 is phenyl, and R^5 , R^7 and R^8 are H, then R^6 cannot be Br.

20 More preferred compounds useful in this invention


25 have the formula:



wherein

R¹ is cyclohexyl,

5



R³ is H or alkyl of 1 to 2 carbon atoms; and/or

R⁴ is CO₂H, a sodium or potassium salt thereof, or

10 CO₂R¹¹; and/or

R⁵ and R⁶ are independently H, halogen or CF₃ provided

that both R⁵ and R⁶ are not hydrogen; and/or

R¹¹ is (CH₂)₂₋₄NR⁹R^{9A}; and/or

R⁹ and R^{9A} are independently alkyl of 1 to 3 carbon

15 atoms, and/or

W and Z are independently H, halogen, alkyl of 1 to 5 carbon atoms or CF₃;

provided that when R¹ is phenyl or phenoxy, and R⁵ is H,

then R⁶ cannot be Br; and that when R¹ is cyclohexyl

20 and R³ is H, R⁶ must be Cl or F.

Specifically preferred compounds useful in this invention are:

- (1) 2-(1,1'-biphenyl-4-yl)-5-chloro-3-methyl-4-quinoline carboxylic acid, sodium or potassium salt

25

- (2) 2-(1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid, sodium or potassium salt
- 5 (3) 6-fluoro-3-methyl-2-(4-phenoxyphenyl)-4-quinoline carboxylic acid, sodium or potassium salt
- (4) 2-(4'-bromo-1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid, sodium or potassium salt
- 10 (5) 2-(2'-fluoro-1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid, sodium or potassium salt.

Detailed Description of the Invention

15 The 4-quinoline-carboxylic acid derivatives useful in this invention are described in and prepared by methods set forth in U.S. Patents 4,680,299 and 4,847,381, the disclosure, synthesis, and synthesis examples of which are hereby incorporated by reference.

20 The isolation of the FK506 natural product is described in European Patent Application publication number 240,773, published 10/14/87 and the chemical synthesis of FK506 is described in Jones et al. (1989) J. Am. Chem. Soc. 111:1157-1159.

25 The preparation of azathioprine is described in U.S. Patent 3,056,785 issued to Burroughs Wellcome. Azathioprine is available as Imuran[®], for which the product information, including dosage and administration, is given in Physicians' Desk Reference

30 44th Edition, 1990, pp 777-778.

The preparation of cyclosporin A is described in U.S. Patent 4,117,118 issued to Sandoz. Cyclosporin A is available as Sandimmune[®], for which the product information, including dosage and information, is given

in Physicians' Desk Reference 44th Edition, 1990, pp 1950-1952.

The preparation of prednisone is described in U.S. Patents 2,897,216 and 3,134,718 issued to Schering.

5 Prednisone is available commercially from several manufacturers as are other corticosteroids (see generally, Physicians' Desk Reference, supra).

Murine monoclonal antibody to the human T3 antigen (herein referred to as OKT3) is available as Orthoclone
10 OKT®3, for which the product information, including dosage and administration and references to methods of preparation, is given in PDR, 1990, pp 1553-1554.

The preparation of mycophenolic acid is described in British patents 1,157,099; 1,157,100; and 1,158,387
15 issued to ICI.

15-dioxyspergualin is a derivative of spergualin discovered in culture filtrates of the bacterial strain BGM162-aFZ as reported in Ochiai, T., Hori, S., Nakajimak, et al. Prolongation of Rat Heart Allograft
20 Survival by 15-desoxyspergualin, 5. Antibiot (Tokyo) 1987; 40:249.

Mizoribine is described in U.S. Patent 3,888,843 issued to Toyo Jozo.

Misoprostol, a prostaglandin (PGE1) analog, is
25 described in U.S. Patent 3,965,143 assigned to Searle and U.S. Patent 4,132,738 assigned to Miles.

Rapamycin is described in U.S. Patents 4,650,803; 4,316,885; 4,885,171; 3,993,749 and 3,929,992, all assigned to Ayerst.

30 Antibodies to the IL-2 receptor protein are described in U.S. Patents 4,578,335 and 4,845,198 (Immunex) and USSN 7/341,361 and U.S. Patent 4,892,827 issued to Pastan et al.

UtilityContact Sensitivity Response to DNFB in Mice

Balb/c female mice (\approx 20 g, Charles River) were sensitized on the shaved abdomen with 25 μ l of 0.5% 2,4-dinitrofluorobenzene (DNFB, Eastman Kodak Co.) in a vehicle of 4:1 acetone:olive oil on days 0 and 1. Mice were ear challenged with 20 μ l of 0.2% DNFB in a vehicle of 4:1 acetone:olive oil on day 5. An identical segment of the ear was measured immediately before challenge and 24 hours later with an engineer's micrometer. Ear swelling was expressed as the difference in ear thickness before and after challenge in units of 10^{-4} inches \pm SEM. Percent suppression was calculated as:

$$\% \text{ Suppression} = 1 - \frac{\text{compound treated-negative control}}{\text{positive control-negative control}} \times 100$$

Compounds were administered orally from day 2 through day 6 and were prepared in 0.25% Methocel[®] (Dow Chemical Co.). Control animals received only vehicle (0.25% Methocel[®]). Negative controls were not sensitized on days 0 and 1 but were ear challenged on day 5. Ten mice were used per group. Results with compounds of invention are shown in Tables 1-4.

Results of the biological tests described below establish that the combination of (1) a 4-quinoline-carboxylic acid derivative and (2) at least one additional immunosuppressive agent, such as cyclosporin A, azathioprine or prednisone, has the effect of suppressing or inhibiting the contact sensitivity response to 2,4-dinitrofluorobenzene (DNFB) in mice.

Contact sensitivity to DNFB is a form of delayed-type hypersensitivity which has been extensively studied to gain an understanding of the regulation of immunologic processes (Claman et al. (1980), Immunological Rev. 50:105-132). This reaction is

- mediated by T lymphocytes that become sensitized to antigen by proliferating and developing into mature effector cells (Claman et al. (1980), Immunological Rev. 50:105-132). This cell-mediated immune response (T-cell mediated immunity) is central to many disease states such as organ transplantation rejection and graft versus host disease (Benacerraf and Unanue (1979), Textbook of Immunology, Williams & Wilkins Co.; Eisen (1980), Immunology, An Introduction to Molecular and Cellular Principles of the Immune Responses, Harper & Row, Inc.; Loveland and McKenzie (1982), Immunology, 46:313-320; Gallin, et al. (1988), Inflammation, Basic Principles and Clinical Correlates, Raven Press).

- A representative 4-quinoline-carboxylic acid derivative, 2-(2'-fluoro-1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid (Example 28, U.S. Patent Number 4,680,299; hereinafter referred to as Example 28), was tested individually (Table 1) as was cyclosporin A (Table 1) and in combination with cyclosporin A in the DNFB contact sensitivity model (Table 2).

Table 1

		Ear ^a			
	<u>Treatment</u>	<u>Dose</u> <u>(mg/kg)</u>	<u>Swelling</u> <u>(units ± SEM)</u>	<u>± Sup-</u> <u>pression</u>	<u>ED50 mg/kg</u>
5	Negative	Vehicle	0.74±0.52	-	-
	Positive	Vehicle	74.11±3.78	0	-
10	Cyclosporin A	2.0	56.15±3.74	24.48	70.00
		10.0	66.58±3.75	10.27	
		50.0	47.90±3.76	35.72	
		100.0	7.80±2.04	90.37	
15	Example 28	0.4	51.95±2.33	30.20	0.95
		2.0	25.61±3.39	66.10	
		10.0	6.40±1.09	92.28	
		20.0	4.75±1.20	94.53	

20

^a Increase in ear thickness from day 5 to day 6,
unit = 10⁻⁴ inches

Table 2

Immunosuppressive Effect (% Suppression of Ear Swelling)
of the Combination of Example 28 and Cyclosporin A in a
Mouse DFNB Contact Sensitivity Model

5	Ex. 28	Cyclosporin A (mg/kg)				
	(mg/kg)	0	6.3	12.5	25.0	30.0
	0	0	-9.9±7.2	-4.0±2.6	0.1±3.5	4.2±4.9
	10	0.1	6.5±2.9	7.4±13.2	1.5±8.0	8.7±9.6
		0.3	7.3±2.4	10.2±17.6	15.2±7.5	3.8±7.1
		1.0	31.4±3.3	51.9±4.6	62.1±3.4	77.0±8.5
		3.0	91.9±2.3	97.0±1.4	99.1±3.8	95.7±3.2
		10.0	93.9±2.4	---	---	91.5±8.0
	15	Ex. 28				
		ED50	1.4	1.0	0.8	0.7
		(mg/kg)				
20	Ex. 28	Cyclosporin A			Cyclosporin A	
	(mg/kg)	50.0	70.0	100.0	ED50	
					(mg/kg)	
	0	33.6±5.7	68.9±1.2	92.5±2.0	60.0	
	0.1	28.8±11.4	---	---	>50.0	
	25	0.3	44.7±7.7	66.7±3.6	---	52.0
		1.0	90.8±1.7	94.6±4.0	---	< 6.3
		3.0	93.9±4.5	92.0±6.2	---	< 6.3
		10.0	88.3±9.0	86.5±7.1	---	< 6.3
	30	Ex. 28				
		ED50	0.4	<0.3		
		(mg/kg)				

As shown in Table 2, test results with mice
receiving the combination treatment of Example 28 and
cyclosporin A (CSA) demonstrate that the

coadministration of suboptimal doses of each agent (Example 28 at 1.0 mg/kg and CSA at 30.0 mg/kg) produced strong suppression of this T-cell-mediated immune response. These data demonstrate a remarkable synergistic effect of the two compounds using this combination therapy. Whereas Example 28 alone at 1.0 mg/kg results in a 31.4% suppression of this immune response and cyclosporin A alone at 30.0 mg/kg results in only a 4.2% suppression of immune response, when these agents are administered in combination at these suboptimal doses (1.0 mg/kg Ex. 28 and 30 mg/kg CSA), an 84.6% suppression of the immune response was observed. Thus, the immunosuppressive effect of the agents used in combination greatly exceeds the simple addition of the effect of each agent used alone. As shown in Table 2, treatment with Example 28 at a dose of 1.0 mg/kg reduced the effective ED₅₀ of cyclosporin A from 60 mg/kg to <6.3 mg/kg.

A similar synergism was observed when Example 28 was administered in combination with azathioprine at suboptimal doses (Table 3). Thus, Example 28 at 1.0 mg/kg and azathioprine at 12.5 mg/kg in combination resulted in a 62.5% immunosuppression, whereas when used individually these agents at these suboptimal doses only resulted in 31.4% suppression and 15.9% suppression, respectively.

Table 3

Immunosuppressive Effect (% Suppression of Ear Swelling)
of the Combination of Example 28 and Azathioprine in a
Mouse DFNB Contact Sensitivity Model

5	Ex. 28	Azathioprine (mg/kg)			
	(mg/kg)	0	6.3	12.5	25.0
	0	0	2.5±4.2	15.9±3.3	29.7±4.1
	10	0.1	6.5±2.8	11.5±10.2	20.9±6.1
		0.3	7.3±2.4	1.7±7.7	25.9±7.5
		1.0	31.4±3.3	38.4±3.8	62.4±3.8
		3.0	91.9±2.3	94.0±2.8	96.3±2.7
		10.0	93.9±2.4	---	100.1±2.2
	15	Ex. 28			
		ED50	1.4	1.3	0.8
20	Ex. 28	Azathioprine (mg/kg)		Azathioprine	
	(mg/kg)	50.0	100.0	ED50	
	0	72.1±3.1	97.7±1.6	(mg/kg)	38.0
	0.1	78.5±3.0	---		42.0
	25	0.3	72.0±3.7	101.2±0.5	43.0
		1.0	90.3±3.4	99.6±3.3	7.5
		3.0	99.2±4.3	100.5±0.2	<6.3
		10.0	101.4±4.4	101.5±3.8	<6.3
	30	Ex. 28			
		ED50	<0.1	<0.1	
35		(mg/kg)			

As the data in Tables 2 and 3 suggest, a lower dose
of each immunosuppressive agent can be used in the
combination of the present invention while maintaining

an effective level of suppression. The dose of the combination to be administered could be reduced on a single dose basis or on a cumulative basis. For example, the number of days on which dosing occurs may be reduced. This is shown in Table 4, wherein dosing was only at day 0 and 1, whereas in previous examples, dosing was on day 0-6.

Table 4

Immunosuppressive Effect (% Suppression of Ear Swelling) of the Combination of Example 28 and Cyclosporin A in a Mouse DNFB Contact Sensitivity Model

Group ^a /Treatment	mg/kg	Δ Ear Swelling ^b	% Sup- pression
Negative Control ^c	-	5.8 \pm 0.6	-
Positive Control	-	68.6 \pm 3.7	-
Ex. 28	20.0	59.6 \pm 2.5	14.4
CSA	100.0	71.8 \pm 3.2	-5.1
Ex. 28 & CSA	20.0 + 100.0	10.8 \pm 1.5	92.0

^a Group of 10 Balb/c female mice were sensitized on days 0 & 1 (0.5% DNFB).

^b All mice were ear challenged on day 5 (0.2% DNFB) and measured on day 6.

^c Nonsensitized control.

The data show that the combination when dosed on days 0 and 1 yielded 92% suppression, whereas cyclosporin A used alone at 100 mg/kg (its optimal dose when administered on days 0-6), when dosed on day 0 and 1 was ineffective (-5.1% suppression).

Current recommended therapy for the prevention of organ transplantation rejection and related disorders,

such as graft versus host disease, involves patient treatment with cyclosporin A and adjunctive therapy with corticosteroids and other immunosuppressive drugs. The present results show that 4-quinoline-carboxylic acid derivatives, such as 2-(2'-fluoro-1,1'-biphenyl-4-yl)-6-(fluoro-3-methyl-4-quinoline carboxylic acid, will be useful as drugs to be used in conjunction with current clinical regimens in synergistically effective amounts. The clinically observed toxicities associated with cyclosporin A (nephrotoxicity), and AZA (hepatotoxicity), have not been seen with the 4-quinoline-carboxylic acid derivatives useful in this invention. Moreover, the 4-quinoline-carboxylic acids derivatives useful in this invention, such as Example 28, have a unique mechanism of action, i.e., inhibition of dihydroorotate dehydrogenase and consequent inhibition of pyrimidine nucleotide biosynthesis (Chen et al. (1986) Cancer Research, 46:5014-5019). By combining a 4-quinoline-carboxylic acid compound of the present invention with another immunosuppressive agent, lower doses of each agent or less frequent administration of each agent, can be used with comparable effectiveness and lower potential for toxic side effects.

The present results show that Example 28 and related 4-quinoline-carboxylic derivatives should be useful when included in combination with other drugs used in current regimens of drug therapy for the prevention of organ transplantation rejection and related disorders (Jacobs and Elgin (1984) "Cyclosporin A, Current Status, Including the Cape Town Experience", in Immune Modulation Agents and Their Mechanisms, pp 191-228, Transplantation and Clinical Immunology, Volume XX Combined Immunosuppressive Therapy in Transplantation ISBN 0-444-81068-4, 1989.

DOSAGE FORMS

The immunosuppressive compounds (active ingredients) of this invention, including at least one 4-quinoline-carboxylic acid compound and at least one immunosuppressive agent preferably selected from the group consisting of, but not limited to: cyclosporin A, azathioprine, corticosteroids such as prednisone, OKT3, FK506, mycophenolic acid or the morpholinethylester thereof, 15-dioxyspergualin, rapamycin, mizoribine, misoprostol and anti-interleukin-2 (IL-2) receptor antibodies, can be administered to treat immunomodulatory disorders and inflammatory diseases and particularly to prevent/treat organ transplantation rejection, graft versus host disease, psoriasis and related disorders, by any means that produces contact of the active ingredient(s) with the agent's site of action in the body of a mammal. The combination therapy can be administered by any conventional means available for use in conjunction with pharmaceuticals; either as individual therapeutically active ingredients or in a combination of therapeutically active ingredients. They can be administered alone, but are generally administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage administered will be an immunosuppressive effective amount of active ingredient(s) and will, of course, vary depending upon known factors such as the pharmacodynamic characteristics of the particular active ingredient, and its mode and route of administration; age, health, and weight of the recipient; nature and extent of symptoms; kind of concurrent treatment, frequency of treatment, and the effect desired. Usually a daily dosage of active ingredient can be about 0.1 to 400 milligrams per


kilogram of body weight. Ordinarily 1 to 100, and preferably 0.1 to 10 milligrams per kilogram per day is effective to obtain desired results.

Pharmaceutical compositions containing the
5 immunosuppressive agents of this invention alone or in combination, can be made by those skilled in the art, using references such as Remington's Pharmaceutical Sciences, A. Osol, a standard reference text in this field. The active ingredients can be administered
10 (alone or in combination) orally in solid dosage forms such as elixirs, syrups, and suspensions, or can also be administered parenterally in sterile liquid dosage forms.

These dosage forms can generally be used when the
15 compounds of this invention are administered stepwise in conjunction with another therapeutic agent. When the drugs are administered in physical combination, the dosage form and administration route should be selected for compatibility with both drugs.


WHAT IS CLAIMED IS:

1. A method of treating organ transplantation rejection in a mammal comprising administering to the mammal a synergistically effective amount of a combination of (a) at least one immunosuppressive agent and (b) at least one compound of the formula:




- 10 or a pharmaceutically acceptable salt thereof wherein:


R is



or




- 15 R¹ is CH₃CH₂(CH₃)CH, alkyl of 5 to 12 carbon atoms, cyclohexyl,



when R is S(O)_mR¹, then

- 20 R¹ can be in addition alkyl of 3 to 4 carbon atoms;
R² is

21




- R^3 is H, alkoxy of 1 to 3 carbon atoms, alkyl of 1 to 2 carbon atoms hydroxy or alkanoyloxy of 2 to 6 carbon atoms;
- when R^3 is hydroxy or alkanoyloxy of 2 to 6 carbon atoms, R^1 can be in addition trifluoromethyl, halogen, hydroxy, alkyl of 1 to 6 carbon atoms and cycloalkyl of 1 to 6 carbon atoms;
- R^4 is CO_2H or CO_2R^{11} ;
- R^5 , R^6 , R^7 and R^8 are independently H, F, Cl, Br, I, CH_3 , CF_3 , SCH_3 or alkyl of 1 to 3 carbon atoms, at least two of R^5 , R^6 , R^7 and R^8 being H;
- R^9 and R^{9A} are independently H or alkyl of 1 to 3 carbon atoms;
- R^{11} is $(\text{CH}_2)_2\text{--}_4\text{NR}^9\text{R}^{9A}$;
- W, Y and Z are independently H, F, Cl, Br, alkyl of 1 to 5 carbon atoms, NO_2 , OH, CF_3 or OCH_3 ; and
- m is 0 or 1;
- with the following provisos:
- (1) R^5 , R^6 and R^7 cannot all be H;
 - (2) when R^4 is $\text{CO}_2\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$, R^6 is CH_2CH_3 , or R^7 is Cl, R^1 cannot be cyclohexyl;
 - (3) when R^1 is cyclohexyl and R^3 is H, R^6 must be Cl or F, but R^6 and R^8 cannot both be Cl; and
 - (4) when R^6 is CH_3 , then R^7 cannot be Cl.
2. The method of Claim 1 wherein the immuno-suppressive agent (a) is selected from the group consisting of: cyclosporin A, azathiopine, OKT3, FK506, mycophenolic acid or the morpholinethylester thereof,

15-dioxyspergualin, rapamycin, mizoribine, misoprostol or an anti-interleukin-2 (IL-2) receptor antibody.

3. The method of Claim 2 wherein the compound has the formula:


5



or a pharmaceutically acceptable salt thereof wherein:


- 10 R^1 is cyclohexyl; phenyl; phenyl substituted with one halogen; alkyl of 1 to 5 carbon atoms or CF_3 ; phenoxy; or phenoxy substituted with one halogen or alkyl of 1 to 5 carbon atoms;
- R^3 is H or alkyl of 1 to 2 carbon atoms;
- 15 R^4 is CO_2H , a sodium or potassium salt thereof; or CO_2R^{11} ;
- R^5 and R^6 are independently H, halogen, CH_3 or CF_3 ;
- R^7 and R^8 are independently H or halogen;
- R^{11} is $(CH_2)_2-4NR^9R^{9A}$; and
- 20 R^9 and R^{9A} are independently alkyl of 1 to 3 carbon atoms;
- provided that R^5 , R^6 and R^7 cannot all be H; when R^1 is cyclohexyl and R^3 is H, R^6 must be Cl or F, but R^6 and R^8 cannot both be Cl; and when R^6 is CH_3 , then
- 25 R^7 cannot be Cl.

4. The method of Claim 2 wherein the compound has the formula:



or a pharmaceutically acceptable salt thereof
wherein:

5 R^1 is cyclohexyl,



- R^3 is H or alkyl of 1 to 2 carbon atoms;
10 R^4 is CO_2H , a sodium or potassium salt thereof, or CO_2R^{11} ;
 R^5 and R^6 are independently H, halogen or CF_3 provided that both R^5 and R^6 are not hydrogen;
 R^{11} is $(\text{CH}_2)_2\text{-}_4\text{NR}^9\text{R}^{9A}$;
15 R^9 and R^{9A} are independently alkyl of 1 to 3 carbon atoms, and
W and Z are independently H, halogen, alkyl of 1 to 5 carbon atoms or CF_3 ;
provided that when R^1 is phenyl or phenoxy, and R^5 is H,
20 then R^6 cannot be Br; and when R^1 is cyclohexyl and R^3 is H, R^6 must be Cl or F.

5. The method of Claim 2 wherein the compound of formula (I) is 2-(1,1'-biphenyl-4-yl)-5-chloro-3-methyl-4-quinoline carboxylic acid, or the sodium or potassium
25 salt thereof.

6. The method of Claim 2 wherein the compound of formula (I) is 2-(1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-

4-quinoline carboxylic acid, or the sodium or potassium salt thereof.

7. The method of Claim 2 wherein the compound of formula (I) is 6-fluoro-3-methyl-2-(4-phenoxyphenyl)-4-quinoline carboxylic acid, the sodium or potassium salt.

8. The method of Claim 2 wherein the compound of formula (I) is 2-(4'-bromo-1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid, sodium or potassium salt.

9. The method of Claim 2 wherein the compound of formula (I) is 2-(2'-fluoro-1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid, sodium or potassium salt.

10. The method of Claim 5 wherein the 4-quinoline-carboxylic acid compound is used in combination with cyclosporin A.

11. The method of Claim 9 wherein the 4-quinoline-carboxylic acid compound is used in combination with cyclosporin A.

12. The method of Claim 5 wherein the 4-quinoline-carboxylic acid compound is used in combination with FK506.

13. The method of Claim 9 wherein the 4-quinoline-carboxylic acid compound is used in combination with FK506.

14. The method of Claim 5 wherein the 4-quinoline-carboxylic acid compound is used in combination with azathioprine.

15. The method of Claim 9 wherein the 4-quinoline-carboxylic acid compound is used in combination with azathioprine.

16. The method of Claim 5 wherein the 4-quinoline-carboxylic acid compound is used in combination with mycophenolic acid or the morpholinethylester thereof.

17. A method of Claim 9 wherein the 4-quinoline-carboxylic acid compound is used in combination with mycophenolic acid or the morpholinethylester thereof.


18. The method of Claim 5 wherein the 4-quinoline-carboxylic acid compound is used in combination with OKT3.

19. The method of Claim 9 wherein the 4-quinoline-carboxylic acid compound is used in combination with OKT3.

20. 20. The method of Claim 5 wherein the 4-quinoline-carboxylic acid compound is used in combination with 15-dioxyspergualin.

21. The method of Claim 9 wherein the 4-quinoline-carboxylic acid compound is used in combination with 15-dioxyspergualin.

22. A method of treating an autoimmune disease graft versus host disease, psoriasis or chronic inflammatory diseases in a mammal comprising administering to the mammal in an amount effective for the treatment of a desired aforesaid disease, a combination of (a) at least one immunosuppressive agent and (b) at least one compound of the formula:




25

or a pharmaceutically acceptable salt, thereof


wherein:

R is




R¹ is CH₃CH₂(CH₃)CH, alkyl of 5 to 12 carbon atoms, cyclohexyl,

5



when R is S(O)_mR¹, then

R¹ can be in addition alkyl of 3-4 carbon atoms;
R² is



10

R³ is H, alkoxy of 1 to 3 carbon atoms, alkyl of 1 to 2 carbon atoms hydroxy or alkanoyloxy of 2 to 6 carbon atoms;

15

when R³ is hydroxy or alkanoyloxy of 2 to 6 carbon atoms, R¹ can be in addition trifluoromethyl, halogen, hydroxy, alkyl of 1 to 6 carbon atoms, and cycloalkyl of 1 to 6 carbon atoms;

R⁴ is CO₂H or CO₂R¹¹;

20

R⁵, R⁶, R⁷ and R⁸ are independently H, F, Cl, Br, I, CH₃, CF₃, SCH₃ or alkyl of 1 to 3 carbon atoms, at least two of R⁵, R⁶, R⁷ and R⁸ being H;

R⁹ and R^{9A} are independently H or alkyl of 1 to 3 carbon atoms;

R¹¹ is (CH₂)₂₋₄NR⁹R^{9A};

W, Y and Z are independently H, F, Cl, Br, alkyl of 1 to 5 carbon atoms, NO₂, OH, CF₃ or OCH₃; and m is 0 or 1;

with the following provisos:

- (1) R⁵, R⁶ and R⁷ cannot all be H;
- (2) when R⁴ is CO₂CH₂CH₂N(CH₃)₂, R⁶ is CH₂CH₃, or R⁷ is Cl, R¹ cannot be cyclohexyl;
- (3) when R¹ is cyclohexyl and R³ is H, R⁶ must be Cl or F, but R⁶ and R⁸ cannot both be Cl; and
- (4) when R⁶ is CH₃, then R⁷ cannot be Cl.

23. The method of Claim 22 wherein the immunosuppressive agent (a) is selected from the group consisting of cyclosporin A, azathioprine, a corticosteroid, OKT3, FK506, mycophenolic acid or the morpholinethylester thereof, 15-dioxyspergualin, rapamycin, mizoribine, misoprostol or an anti-interleukin-2 (IL-2) receptor antibody.

24. The method of Claim 23 wherein the compound of formula (I) is 2-(1,1'-biphenyl-4-yl)-5-chloro-3-methyl-4-quinoline carboxylic acid, or the sodium or potassium salt thereof.

25. The method of Claim 23 wherein the compound of formula (I) is 2-(1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid, or the sodium or potassium salt thereof.

26. The method of Claim 23 wherein the compound of formula (I) is 6-fluoro-3-methyl-2-(4-phenoxyphenyl)-4-quinoline carboxylic acid, or the sodium or potassium salt thereof.

27. The method of Claim 23 wherein the compound of formula (I) is 2-(4'-bromo-1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid, or the sodium or potassium thereof.

5 28. The method of Claim 23 wherein the compound of formula (I) is 2-(2'-fluoro-1,1'-biphenyl-4-yl)-6-fluoro-3-methyl-4-quinoline carboxylic acid, or the sodium or potassium thereof.

10 29. The method of Claim 24 wherein the 4-quinoline-carboxylic acid compound is used in combination with cyclosporin A.

30. The method of Claim 28 wherein the 4-quinoline-carboxylic acid compound is used in combination with cyclosporin A.

15 31. The method of Claim 24 wherein the 4-quinoline-carboxylic acid compound is used in combination with FK506.

20 32. The method of Claim 28 wherein the 4-quinoline-carboxylic acid compound is used in combination with FK506.

33. The method of Claim 24 wherein the 4-quinoline-carboxylic acid compound is used in combination with azathioprine.

25 34. The method of Claim 28 wherein the 4-quinoline-carboxylic acid compound is used in combination with azathioprine.

30 35. The method of Claim 24 wherein the 4-quinoline-carboxylic acid compound is used in combination with mycophenolic acid or the morpholinethylester thereof.

36. The method of Claim 28 wherein the 4-quinoline-carboxylic acid compound is used in combination with mycophenolic acid or the morpholinethylester thereof.

37. The method of Claim 24 wherein the 4-quinoline-carboxylic acid compound is used in combination with OKT3.

38. The method of Claim 28 wherein the 4-quinoline-carboxylic acid compound is used in combination with OKT3.

39. The method of Claim 24 wherein the 4-quinoline-carboxylic acid compound is used in combination with prednisone.

40. The method of Claim 28 wherein the 4-quinoline-carboxylic acid compound is used in combination with prednisone.

41. The method of Claim 24 wherein the 4-quinoline-carboxylic acid compound is used in combination with 15-dioxyspergualin.

42. The method of Claim 28 wherein the 4-quinoline-carboxylic acid compound is used in combination with 15-dioxyspergualin.

43. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and at least one immunosuppressive agent (other than a 4-quinoline-carboxylic acid derivative) and at least one compound of Formula I, as defined in Claim 1, as a combined preparation permitting the simultaneous or separated sequential use of said immunosuppressive agent and said 4-quinoline-carboxylic acid derivative of Formula I.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US91/03788

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ³

According to International Patent Classification (IPC) or to both National Classification and IPC

IPC(5): A61K 31/47, A61K 37/00

II. FIELDS SEARCHED

Minimum Documentation Searched ⁴

Classification System

Classification Symbols

U.S. CL. 514-15, 514-311, 514-312

Documentation Searched other than Minimum Documentation
to the extent that such Documents are Included in the Fields Searched ⁵

III. DOCUMENTS CONSIDERED TO BE RELEVANT ^{1,2}

Category ⁶	Citation of Document, ^{1,2} with indication, where appropriate, of the relevant passages ^{1,7}	Relevant to Claim No. ^{1,8}
Y	US, A 3,574,840 (RIVIERE) 13 APRIL 1971	1-11, 22-30&43
Y	US, A 4,117,135 (SCHRODER) 26 SEPTEMBER 1978	1-11, 22-30&43
Y	US, A 4,861,783 (ACKERMAN) 29 JULY 1989	1-11, 22-30&43
Y	US, A 4,661,499 (YOUNG) 28 APRIL 1987	1-11, 22-30&43
Y	US, A 4,680,,299 (HESSON) 14 JULY 1987	1-11, 22-30&43
Y	US, A 3,973,022 (GOSCHUE) 03 AUGUST 1976	1-11, 22-30&43
Y	US, A 4,929,626 (MOHRS) 29 MAY 1990	1-11, 22-30&43
Y	US, A 4,851,409 (YOUNG) 25 JULY 1989	1-11, 22-30&43
Y	EP, A 0,213,151 (LARSON) 05 AUGUST 1987	1-11, 22-30&43
Y	Chemical Abstract, volume 111, No. 3, 1989, 17327V. (GERDIN), "Rat cardiac allografts protected with cyclosporin a are rejected in the presence of LS-2616 (LINOMIDE).	1-11, 22-30&43

⁹ Special categories of cited documents: ¹³

"A" document defining the general state of the art which is not
considered to be of particular relevance

"E" earlier document but published on or after the international
filing date

"L" document which may throw doubts on priority claim(s) or
which is cited to establish the publication date of another
citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or
other means

"P" document published prior to the international filing date but
later than the priority date claimed

"T" later document published after the international filing date
or priority date and not in conflict with the application but
cited to understand the principle or theory underlying the
invention

"X" document of particular relevance: the claimed invention
cannot be considered novel or cannot be considered to
involve an inventive step

"Y" document of particular relevance: the claimed invention
cannot be considered to involve an inventive step when the
document is combined with one or more other such docu-
ments, such combination being obvious to a person skilled
in the art.

"Δ" document member of the same patent family

IV. CERTIFICATION

Date of the Actual Completion of the International Search ²

10 JULY 1991

Date of Mailing of this International Search Report ²

19 AUG 1991

International Searching Authority ¹

ISA/US

Signature of Authorized Officer ²⁰

STANLEY J. FRIEDMAN